

A Novel Ameliorant Biochar of Areca Nuts Skin and Sago Bark Waste for Increasing Soil Chemical Fertility

Febrianti Rosalina^{a,*}, Ihsan Febriadi^b

^a Department of Agrotechnology, Universitas Muhammadiyah Sorong, Sorong, 98412, Indonesia

^b Department of Agroforestry, Universitas Muhammadiyah Sorong, Sorong, 98412, Indonesia

Corresponding author: *febriantirosalina@um-sorong.ac.id

Abstract— The waste of areca nuts skin and sago bark are alternatives of ameliorant material that can be used to support soil quality improvement. This research aims to study the potential of areca nuts and sago bark to be produced as biochar and to identify the chemical components they contain. The study employed a Completely Randomized Designed involving six treatments repeated three times. The observation parameters of soil chemical analysis, including the measurement of pH H₂O (pH meter); Total-N (Kjeldahl); Available-P (Bray I); Exchangeable Cations (NH₄OAC pH 7.0); and exchangeable-Al (KCl 1N). Data obtained from observation were statistically analyzed through the ANOVA method, and if they had a significant effect, the analysis was continued with Duncan's Multiple Range Test (DMRT) at the level of $\alpha=5\%$. Findings showed that the administration of biochar can improve soil quality. Based on the analysis, biochar from sago bark waste brings a significant effect on increases of pH (5.26), available-P (12.99 cmol⁺kg⁻¹), Exchangeable-K (2.16 cmol⁺kg⁻¹), Exchangeable-Na (0.21 cmol⁺kg⁻¹), Exchangeable-Ca (4.83 cmol⁺kg⁻¹), Exchangeable-Mg (3.80 cmol⁺kg⁻¹), and the reduction of soil Exchangeable-Al (6.65 cmol⁺kg⁻¹). In comparison, biochar from betel nut peel waste has a significant effect on Total-N (0.15%), C-organic (1.66%), and CEC (23.38 cmol⁺kg⁻¹). The results of this study will be the basis for further research in utilizing agricultural waste so that the right combination of fertilizers is obtained to increase soil fertility and will indirectly increase crop production, especially in West Papua.

Keywords— Ameliorant; areca nuts skin; biochar; sago bark; waste.

Manuscript received 19 Sep. 2022; revised 9 Jan. 2023; accepted 15 Feb. 2023. Date of publication 30 Jun. 2023.
IJASEIT is licensed under a Creative Commons Attribution-Share Alike 4.0 International License.



I. INTRODUCTION

Indonesia is rich in natural resources. Among many areas in Indonesia, Papua has abundant natural resources. One of them, in the agricultural sector, is the commodities of Areca and Sago. Data obtained from the Indonesian Plantation Statistic indicated that the area of sago farms in Papua and West Papua was 37,017 ha, with an estimated production of 29,818 tons. While the area of areca nuts plantations in Papua Province is 1,653 ha, with a total production of 350 tons. Areca nuts and sago are typical Papuan fruits, and Papuan societies widely use Areca nuts as stimulants mixed with betel and lime. Besides that, areca nut is also used for food, industrial raw materials such as fabric dyes, medicine, and antioxidants [1], [2].

In comparison, sago is the staple food of local indigenous people and is used as an alternative material for boats [3], [4]. Based on an earlier study, sago waste can be used as a vegetable herbicide [5], [6]. Sago waste can be used for

feeding animals, plant medium, and material for bio-foam [7]–[10]. Besides, Papua local sago extraction liquid waste is used as media to grow microalgae for biodiesel production [11], [12]. Many studies have been conducted to investigate the use of areca nut and sago waste, yet the agricultural sector is still very limitedly implemented. Unfortunately, the study about using areca nut waste to improve soil quality is limited. Processing areca nut and sago will result in solid, liquid, and gaseous waste. They produce two types of solid waste: bark and dregs, which are 75% in total [13].

The abundant amount of areca nut and sago bark waste has not been optimally used in Papuan societies, and it can turn into a serious environmental problem if not handled appropriately. Therefore, waste management is urgently needed. It is considered that most areas of Papua, especially West Papua, are dry and infertile, and it is important to add nutrients and manage them appropriately to utilize them. One of the methods to improve the quality of farmland is by utilizing waste of areca nut skin and sago bark.

The alternative utilization of waste of areca nut skin and sago bark is processing it as biochar as a soil ameliorant. Biochar is a soil ameliorant produced from burning organic matter (pyrolysis) with limited oxygen conditions [14]. The application of biochar can increase acid soil pH [15], [16] increase soil Cation Exchange Capacity (CEC), and provide the nutrition of N, P, and K [17]–[20]. Biochar maintains the soil moisture so that the capacity keeps the high level of water [21] and repairs soil contaminated with heavy metals like Cu, Pb, Ni, and Cd [22]–[25]. Besides that, biochar contained in the soil can boost growth and support the plants to absorb nutrients [26]–[29]. Sago bark is recommended to be used as biochar because the yield of charcoal from sago bark is higher than from sago pith [30]. It is because sago bark contains more lignin which produces more carbon. Thus the higher the lignin, the more the yield [31], [32]. The benefits of areca nut waste are that it can improve soil quality by acting as an ameliorant and binding heavy metals in solution [33]–[35].

Based on the background, the study aims to identify the potential of areca nut and sago bark waste as biochar and measure the decomposition of nutrient content for increasing soil chemical fertility. The use of waste of areca nut and sago bark is expected to provide initial data that can be used for further wider-scale studies. It can contribute to improving soil

structure and quality to indirectly increase agricultural production in the eastern part of Indonesia, especially Sorong, West Papua.

II. MATERIALS AND METHOD

The study employed a completely randomized design consisting of 9 treatments including LP0 (0 ton ha⁻¹ areca nut biochar), LP1 (40 ton ha⁻¹ areca nut biochar), LP2 (80 ton ha⁻¹ areca nut biochar), LS0 (0 ton ha⁻¹ sago biochar), LS1 (40 ton ha⁻¹ sago biochar), LS2 (80 ton ha⁻¹ sago biochar), LPS0 (0 ton ha⁻¹ areca nut + sago biochar), LPS1 (40 ton ha⁻¹ areca nut + sago biochar), LPS2 (80 ton ha⁻¹ areca nut + sago biochar), and each treatment was repeated three times.

The main material of the biochar was areca nut skin and sago bark, which had been dried to reduce the water content. After that, the stove was made by digging the soil to make a hole like half of the ball (diameter: 1.5 m; depth: 50 cm). We used a chimney (diameter: 30-35 cm) to supply oxygen. The main material is put into the chimney, and the combustion is started from inside the chimney using flammable material. Biochar produced from the combustion was then weighted according to the doses of ameliorant needed in each treatment.




Fig. 1 Flowchart of research method

Soil for incubation media was taken from 0-20 cm depth in different locations and composited, air-dried, and filtered using a sieve with a hole diameter of 5 mm. After that, 0.5 kg (air dry weight) of it was taken and put into an incubation

container and mixed with biochar based on the doses of each treatment. The incubation media is added with basic fertilizer: 0.06-gram pot⁻¹ (100 kg ha⁻¹ SP-36), 0.04-gram pot⁻¹ (100 kg ha⁻¹ KCl), and 0.14-gram pot⁻¹ (300 kg ha⁻¹ Urea). The

incubation container was kept in the laboratory, and the soil condition was retained under field capacity conditions. After six weeks of incubation, the soil sample was taken and analyzed in the laboratory [36].


Chemical analysis of incubation media included pH H₂O (pH meter); N-total (Kjeldahl); available-P (Bray I); exchangeable-K, exchangeable-Na, exchangeable-Ca, exchangeable-Mg and CEC (NH₄OAC pH 7.0); exchangeable-Al (KCl 1N). Data obtained from observation were statistically analyzed through the ANOVA method, and if they had a significant effect, the analysis was continued with Duncan's Multiple Range Test (DMRT) at the level of $\alpha=5\%$. The flowchart methodology of this research is in Fig. 1

III. RESULT AND DISCUSSION

The effect of sago bark and areca nut skin biochar on soil's chemical characteristics after incubation for six weeks is presented in Figures 1-5 and Table 1. Results of variance analysis show a real influence of ameliorant materials (biochar) on soil's pH, Organic-C, Total-N, available-P, exchangeable-Na, exchangeable-Ca, exchangeable-K, exchangeable-Mg, CEC, exchangeable-Al, and exchangeable-H.

A. Soil Reaction (pH)

Based on the statistical analysis as presented in Figure 1, it is identified that biochar increases the soil pH. Further analysis shows that LS2 had no significant difference from LPS2 but had a significant difference with other treatments. Based on the graph trend (Fig. 2), the soil pH increased by 4.10 with an average increase of 4.34 – 5.26



Numbers shown in the bar chart followed by similar letters show no significant difference on the 5% test level using Duncan's Multiple Range Test (DMRT).


Fig. 2 The influence of biochar on soil pH

The increase in soil active pH score (pH H₂O) from 4.01 to 5.26 was due to the high pH of the main material (pH 5-6). High pH shows that biochar that is used can be applied as an ameliorant for the soil. Biochar is more appropriate to be used as an amendment (ameliorant) for soil and as an alternative to organic fertilizers [37], [38]. The contribution of the biochar OH also causes an increase in pH. The measurement of the soil pH (pH H₂O) shows that the concentration of H⁺ in the soil solution corresponds to the real environmental condition that is higher than the pH of potential soil (pH KCl or pH CaCl₂) [39]. In this case, the active pH expresses the concentration of H⁺ in the soil solution, and the potential pH expresses the concentration of H⁺ in colloidal soil. Previous studies strengthen that the alkaline nature of biochar is formed from H⁺ ions which can be exchanged with the surrounding

soil leading to the increase of soil pH from applying the biochar [40], [41]. The combination of ameliorant materials that is applied can increase the soil pH and reduce the use of lime for acid soil [42]–[45]

B. Organic Carbon and Total Nitrogen


The statistical analysis shows that adding biochar to the soil can increase the Organic-C content. Based on further analysis, it was identified that LP2 treatment shows a real difference from other treatments (Fig. 3).



Numbers shown in the bar chart followed by similar letters show no significant difference on the 5% test level using Duncan's Multiple Range Test (DMRT).

Fig. 3 The influence of biochar on soil Organic-C content

Biochar can store carbon well, is more resistant to oxidation, and can be more stable in the soil [46], [47]. As per further analysis, it was identified that the treatment of LP2 was different from other treatments. LP2 treatment is applied to areca nut peel waste. Areca nut waste (skin) contains cellulose, water, and ash with a rate of cellulose of around 70,2 % that is suspected can act as the source of carbon for the production of active charcoal [48], [49]. It possibly leads to the LP2 treatment being better than others. However, based on Figure 2, it is identified that the Organic-C of the soil increases based on the dose of each main material used. Sago waste biochar can increase the soil's Organic-C, pH, and CEC [50]–[52]. It confirms [53] that biochar made from sago waste can increase the rate of carbon. Fig. 4 shows the Total-N content of the soil that is inversely proportional with the Organic-C it produces (Figure 2), except soil added with ameliorant produced from areca nut + sago bark waste (LPS0, LPS1, LPS2).



Numbers shown in the bar chart followed by similar letters show no significant difference on the 5% test level using Duncan's Multiple Range Test (DMRT).


Fig. 4 The influence of biochar on soil Total Nitrogen

Fig. 4 shows that the Total-N of the soil was inversely proportional with the Organic-C it produced (Fig. 2), except for the treatment using areca nut-sago bark waste (LPS0,

LPS1, LPS2). It is possible because of the contribution of C from biochar and the increase of microorganism activities. The availability of a higher amount of C stimulates the activities of the microbes, leads to the increase of N demand, pushes immobilization, and recycles NO_3^- , reducing the availability of N [54], [55].

C. Available Phosphor

Based on Figure 5, the more main material used, the higher the available-P of the soil. Among all treatments, the use of sago bark waste shows the best result, although based on the trend (Figure 4), there was a drastic increase from treatments LS0 to LS1. It was also proved by the statistical analysis showing that LS2 treatment shows a striking difference compared to others.



Numbers shown in the bar chart followed by similar letters show no significant difference on the 5% test level using Duncan's Multiple Range Test (DMRT).

Fig. 5 The influence of biochar on available-P of soil

Among all treatments, the one using sago bark waste showed the best result, although based on the trend (Fig. 5),

there was a dramatic increase from LS0 to LS1 treatments. It was also proved by the statistical analysis showing that LS2 treatment was different from others. The administration of biochar can reduce the Al solubility and increase the soil pH so that the P binding can be avoided and the P available in the soil can increase. It follows that the solubility of metal ions such as Al and Fe can reduce the adsorption of P and available-P in the forms of PO_4^{3-} , HPO_4^{2-} , and H_2PO_4^- [56], [57]. Also, adding biochar indicates the increase of soil available-P [56]. Because biochar can increase the soil enzymatic activities involved in C, N, and P cycles [57].

D. Exchangeable Cations

Based on the data presented in Table I, in general, treatments showing the best results were the ones applying the highest doses (LP2, LS2, and LPS2). The oxidation of C aromatic and the formation of the carboxyl group is believed to be the main reason for the increase in CEC [54]. While biochar has a high rate of cation exchange because of the big surface area capacity, leading to an increase in soil pH and water holding capacity, as well as an affinity for plant micro and macronutrients [58]. It confirms that applying biochar as a soil improvement agent on suboptimal and degraded land, besides increasing the soil pH, Total-N, and available-P, can also improve the soil's cation exchange capacity [59], [60]. In this case, it increases exchangeable-K, exchangeable-Na, exchangeable-Ca, and exchangeable-Mg. Applying organic material as an ameliorant can improve the quality of soil's chemical properties [61], [62]. Besides improving the soil pH, biochar can also reduce the leaching of nutrients in the growing media [63].

TABLE I
INFLUENCES OF BIOCHAR ON CATIONS THAT CAN BE EXCHANGE


Treatments	Exchangeable Cations ($\text{cmol}^+\text{kg}^{-1}$)				CEC ($\text{cmol}^+\text{kg}^{-1}$)
	Exc-K	Exc-Na	Exc-Ca	Exc-Mg	
LP0	0.55 ^a	0.14 ^a	3.50 ^a	3.60 ^{cd}	21.86 ^b
LP1	1.16 ^{de}	0.16 ^{ab}	3.98 ^b	3.50 ^{bc}	23.83 ^b
LP2	2.01 ^g	0.21 ^{cd}	4.43 ^c	3.75 ^{de}	22.80 ^b
LS0	0.84 ^{bc}	0.21 ^d	3.74 ^{ab}	3.73 ^{de}	19.41 ^a
LS1	0.98 ^{cd}	0.20 ^{cd}	3.54 ^a	3.76 ^{de}	23.75 ^b
LS2	2.16 ^g	0.21 ^{cd}	4.83 ^d	3.80 ^e	22.65 ^b
LPS0	0.61 ^{ab}	0.14 ^a	3.65 ^{ab}	3.22 ^a	22.32 ^b
LPS1	1.31 ^{ef}	0.18 ^{bc}	3.81 ^{ab}	3.41 ^b	23.42 ^b
LPS2	1.52 ^f	0.19 ^{cd}	3.79 ^{ab}	3.82 ^c	21.86 ^b

Numbers shown in the similar column followed by similar letters show no significant difference on the 5% test level using Duncan's Multiple Range Test (DMRT).

E. Exchangeable Aluminum

The statistical analysis shows that treatments that can significantly affect the reduction of soil exchangeable-Al were the LS2 (biochar from sago bark waste). The treatments indicated that LS2 treatment could reduce the soil exchangeable-Al from 9,11 $\text{cmol}^+\text{kg}^{-1}$ to 6,65 $\text{cmol}^+\text{kg}^{-1}$. Based on the graph in Fig. 6, the higher the doses, the lower the exchangeable-Al in the soil. In this case, administering 80 tons ha^{-1} of sago biochar (LS2) is the best dose to reduce the soil's exchangeable Al. While the interaction of both biochar materials shows a real effect, it is not as good as LS2 treatment in reducing soil's exchangeable Al. It indicated a reaction between materials causing the effect on the parameter reduction. It echoes those who argued that using sago waste biochar could reduce the soil's exchangeable Al [64]. While applying biochar on soil can reduce the

concentration of exchangeable Al due to the reduction of Al, which can be exchanged through adsorption on the biochar surface, which is negatively charged from OH^- [54], [65].



Numbers shown in the bar chart followed by similar letters show no significant difference on 5% test level using Duncan's Multiple Range Test (DMRT)

Fig. 6 The effect of biochar on soil exchangeable-Al

IV. CONCLUSION

The administration of biochar made of areca nut and sago bark waste as ameliorant materials can improve soil quality. It can be seen from the chemical characteristics of the soil after receiving biochar. The analysis shows that the addition of biochar from sago bark waste stimulates the increase of the pH (5.26), available-P (12.99 cmol^+kg^-), exchangeable-K (2.16 cmol^+kg^-), exchangeable-Na (0.21 cmol^+kg^-), exchangeable-Ca (4.83 cmol^+kg^-), exchangeable-Mg (3.80 cmol^+kg^-), and the reduction of the soil exchangeable-Al (6.65 cmol^+kg^-). While biochar from areca nut waste highly influences the rate of total-N (0.15%), Organic-C (1.66%), and CEC (23.38 cmol^+kg^-). The results of this study will be the basis for further research in utilizing agricultural waste so that the right combination of fertilizers is obtained to increase soil fertility and will indirectly increase crop production, especially in West Papua.

ACKNOWLEDGMENT

The Council for Higher Education Research and Development of Muhammadiyah Central Leadership and Universitas Muhammadiyah Sorong have supported this research.

REFERENCES

- [1] F. D. Korwa, J. Husain, T. Titah, and J. Supit, "Evaluasi Kesesuaian Lahan untuk Tanaman Pinang (*Areca catechu*) di Das Remu, Sorong, Papua Barat," *Cocos*, vol. 7, no. 4, pp. 1–8, 2016, doi: 10.35791/cocos.v7i4.12602.
- [2] N. R. Yanti, M. Andika, S. Maulida, Riani, I. Sulaiman, and N. M. Erfiza, "Utilization of areca nut (*Areca chatechu* L.) extract for tannin based colorimetric indicator in smart packaging," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 951, no. 1, pp. 1–7, 2022, doi: 10.1088/1755-1315/951/1/012057.
- [3] F. F. Sidiq, D. Coles, C. Hubbard, B. Clark, and L. J. Frewer, "Sago and the indigenous peoples of Papua, Indonesia: A review," *J. Agric. Appl. Biol.*, vol. 2, no. 2, pp. 138–149, 2021, doi: 10.11594/jaab.02.02.08.
- [4] S. K. Uda, L. Hein, and A. Adventa, "Towards better use of Indonesian peatlands with paludiculture and low-drainage food crops," *Wetl. Ecol. Manag.*, vol. 28, no. 3, pp. 509–526, 2020, doi: 10.1007/s11273-020-09728-x.
- [5] J. Putinella, Y. Nuraini, and B. Prasetya, "Nitrogen released from mixtures of sago pulp waste and *Gliricidia sepium* pruning on a Dystrudept of Central Moluccas and its effect on maize growth of maize," *J. Degrad. Min. Lands Manag.*, vol. 9, no. 2, pp. 3341–3347, 2022, doi: 10.15243/jdmlm.2022.092.3341.
- [6] K. M. T. Sulok *et al.*, "Chemical and biological characteristics of organic amendments produced from selected agro-wastes with potential for sustaining soil health: A laboratory assessment," *Sustain.*, vol. 13, no. 9, pp. 1–15, 2021, doi: 10.3390/su13094919.
- [7] H. P. Wardono, A. Agus, A. Astuti, N. Ngadiyono, and B. Suhartanto, "Potential of sago hampas for ruminants feed," *E3S Web Conf.*, vol. 306, p. 05012, 2021, doi: 10.1051/e3sconf/202130605012.
- [8] N. C. Tiven and T. M. Simanjorang, "Effect of urea in steamed sago waste on rumen fermentation parameters in vitro tested," in *IOP Conference Series: Earth and Environmental Science*, 2021, vol. 883, no. 1, doi: 10.1088/1755-1315/883/1/012054.
- [9] A. L. W. Jampi, S. F. Chin, M. E. Wasli, and C. H. Chia, "Preparation of Cellulose Hydrogel from Sago Pith Waste as a Medium for Seed Germination," *J. Phys. Sci.*, vol. 32, no. 1, pp. 13–26, 2021, doi: 10.21315/JPS2021.32.1.2.
- [10] N. K. Mohamad Fathi, M. F. Mohamad Bukhori, S. M. Abd Aziz Abdullah, R. Wahi, M. A. Zailani, and M. M. Raja Gopal, "Effect OF Sago Bark Biochar Application on *Capsicum annum* L. var. Kulai Growth and Fruit Yield," *Malaysian Appl. Biol.*, vol. 51, no. 3, pp. 127–135, 2022, doi: 10.55230/mabjournal.v51i3.2191.
- [11] K. Thangavelu, P. Sundararaju, N. Srinivasan, and S. Uthandi, "Bioconversion of sago processing wastewater into biodiesel: Optimization of lipid production by an oleaginous yeast, *Candida tropicalis* ASY2 and its transesterification process using response surface methodology," *Microb. Cell Fact.*, vol. 20, no. 1, pp. 1–18, 2021, doi: 10.1186/s12934-021-01655-7.
- [12] R. K. Srivastava, N. P. Shetti, K. R. Reddy, E. E. Kwon, M. N. Nadagouda, and T. M. Aminabhavi, "Biomass utilization and production of biofuels from carbon neutral materials," *Environ. Pollut.*, vol. 276, p. 116731, 2021, doi: 10.1016/j.envpol.2021.116731.
- [13] H. Siruru, W. Syafii, I. N. J. Wistara, G. Pari, and I. Budiman, "Properties of Sago waste charcoal using hydrothermal and pyrolysis carbonization," *Biomass Convers. Biorefinery*, 2020, doi: 10.1007/s13399-020-00983-9.
- [14] D. Spanu, G. Binda, C. Dossi, and D. Monticelli, "Biochar as an alternative sustainable platform for sensing applications: A review," *Microchem. J.*, vol. 159, no. June, p. 105506, 2020, doi: 10.1016/j.microc.2020.105506.
- [15] H. Herviyanti, A. Maulana, S. Prima, A. Aprisal, S. D. Crisna, and A. L. Lita, "Effect of biochar from young coconut waste to improve chemical properties of ultisols and growth coffee [*Coffea arabica* L.] plant seeds," in *IOP Conference Series: Earth and Environmental Science*, 2020, vol. 497, no. 1, doi: 10.1088/1755-1315/497/1/012038.
- [16] S. Wu, Y. Zhang, Q. Tan, X. Sun, W. Wei, and C. Hu, "Biochar is superior to lime in improving acidic soil properties and fruit quality of Satsuma mandarin," *Sci. Total Environ.*, vol. 714, p. 136722, 2020, doi: 10.1016/j.scitotenv.2020.136722.
- [17] A. Abbas *et al.*, "Efficiency of wheat straw biochar in combination with compost and biogas slurry for enhancing nutritional status and productivity of soil and plant," *Plants*, vol. 9, no. 11, pp. 1–19, 2020, doi: 10.3390/plants9111516.
- [18] A. Karimi, A. Moezzi, M. Chorom, and N. Enayatizamir, "Application of Biochar Changed the Status of Nutrients and Biological Activity in a Calcareous Soil," *J. Soil Sci. Plant Nutr.*, vol. 20, no. 2, pp. 450–459, 2020, doi: 10.1007/s42729-019-00129-5.
- [19] M. A. Rodrigues *et al.*, "Nitrogen Use Efficiency and Crop Yield in Four Successive Crops Following Application of Biochar and Zeolites," *J. Soil Sci. Plant Nutr.*, vol. 21, no. 2, pp. 1053–1065, 2021, doi: 10.1007/s42729-021-00421-3.
- [20] W. Zhao, Q. Zhou, Z. Tian, Y. Cui, Y. Liang, and H. Wang, "Apply biochar to ameliorate soda saline-alkali land, improve soil function and increase corn nutrient availability in the Songnen Plain," *Sci. Total Environ.*, vol. 722, p. 137428, 2020, doi: 10.1016/j.scitotenv.2020.137428.
- [21] P. Pokharel and S. X. Chang, "Biochar decreases the efficacy of the nitrification inhibitor nitrapyrin in mitigating nitrous oxide emissions at different soil moisture levels," *J. Environ. Manage.*, vol. 295, no. June, p. 113080, 2021, doi: 10.1016/j.jenvman.2021.113080.
- [22] S. Mansoor *et al.*, "Chemosphere Biochar as a tool for effective management of drought and heavy metal toxicity," *Chemosphere*, vol. 271, p. 129458, 2021, doi: 10.1016/j.chemosphere.2020.129458.
- [23] M. Liang, L. Lu, H. He, J. Li, Z. Zhu, and Y. Zhu, "Applications of biochar and modified biochar in heavy metal contaminated soil: A descriptive review," *Sustain.*, vol. 13, no. 24, pp. 1–18, 2021, doi: 10.3390/su132414041.
- [24] J. Zhang, Z. Tan, and Q. Huang, "Study on principles and mechanisms of new biochar passivation of cadmium in soil," *Biochar*, vol. 3, no. 2, pp. 161–173, 2021, doi: 10.1007/s42773-021-00088-0.
- [25] A. Taraqqi-A-kamal *et al.*, "Biochar remediation of soil: Linking biochar production with function in heavy metal contaminated soils," *Plant, Soil Environ.*, vol. 67, no. 4, pp. 183–201, 2021, doi: 10.17221/544/2020-PSE.
- [26] T. Kocsis, Z. Kotrocó, L. Kardos, and B. Biró, "Optimization of increasing biochar doses with soil–plant–microbial functioning and nutrient uptake of maize," *Environ. Technol. Innov.*, vol. 20, p. 101191, 2020, doi: 10.1016/j.eti.2020.101191.
- [27] A. A. Karim *et al.*, "Enrichment of primary macronutrients in biochar for sustainable agriculture: A review," *Crit. Rev. Environ. Sci. Technol.*, vol. 52, no. 9, pp. 1449–1490, 2022, doi: 10.1080/10643389.2020.1859271.
- [28] N. Khairunnisa *et al.*, "Physicochemical Properties of Sago Bark Biochar and Its Potential As Plant Growth Media (Sifat Fizikokimia Bio-Arang Sisa Kulit Sagu dan Potensinya Sebagai Media Pertumbuhan Tanaman)," *Malaysian J. Anal. Sci.*, vol. 25, no. 4, pp. 622–636, 2021.
- [29] A. A. Rahi *et al.*, "Toxicity of Cadmium and nickel in the context of applied activated carbon biochar for improvement in soil fertility," *Saudi J. Biol. Sci.*, vol. 29, no. 2, pp. 743–750, 2022, doi: 10.1016/j.sjbs.2021.09.035.

- [30] J. Rambli, W. A. Wan Abd Karim Ghani, and M. A. Mohd Salleh, "Characterization of Sago-based Biochar as Potential Feedstock for Solid Fuel," *J. Energy Saf. Technol.*, vol. 1, no. 2, pp. 11–17, 2018, doi: 10.11113/jest.v1n2.16.
- [31] C. Kang, Y. Huang, H. Yang, X. F. Yan, and Z. P. Chen, "A review of carbon dots produced from biomass wastes," *Nanomaterials*, vol. 10, no. 11, pp. 1–24, 2020, doi: 10.3390/nano10112316.
- [32] O. Togibasa, M. Mumfajjah, Y. K. Allo, K. Dahlan, and Y. O. Ansanay, "The effect of chemical activating agent on the properties of activated carbon from sago waste," *Appl. Sci.*, vol. 11, no. 24, 2021, doi: 10.3390/app112411640.
- [33] Y. Istikorini, Nurhafifah, A. P. P. Hartoyo, A. Solikhin, and E. A. Octiaviani, "Effect of plant growth-promoting rhizobacteria and bionanomaterial membrane applications on chemical properties of peat soils," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 959, no. 1, 2022, doi: 10.1088/1755-1315/959/1/012049.
- [34] B. Weerasuk, S. ang Supharoek, K. Grudpan, and K. Ponghong, "Exploiting crude betel nut (*Areca catechu* Linn.) extracted solution as a natural reagent with sequential injection spectrophotometry for iron analysis in rice samples," *J. Iran. Chem. Soc.*, vol. 19, no. 3, pp. 741–751, 2022, doi: 10.1007/s13738-021-02337-2.
- [35] T. M. Novera, M. Tabassum, M. Bardhan, M. A. Islam, and M. A. Islam, "Chemical modification of betel nut husk prepared by sodium hydroxide for methylene blue adsorption," *Appl. Water Sci.*, vol. 11, no. 4, pp. 1–14, 2021, doi: 10.1007/s13201-021-01394-5.
- [36] F. Rosalina, D. Tjahyandari, and D. Darmawan, "The Potential Of Nickel Slag with Humic Substance Addition as Ameliorating Materials on Gajrug Red-Yellow Podzolic," *Sains Tanah - J. Soil Sci. Agroclimatol.*, vol. 15, no. 1, p. 61, 2018, doi: 10.15608/stjssa.v15i1.17814.
- [37] H. Senghie, M. H. Bolhassan, and D. S. Awg-Adeni, "The Effects of Sago (Metroxylon sago) Bark and Frond Waste as Substrates on the Growth and Yield of Grey Oyster Mushrooms (*Pleurotus sajor-caju*)," *Pertanika J. Trop. Agric. Sci.*, vol. 44, no. 2, pp. 307–316, 2021, doi: 10.47836/PJTAS.44.2.04.
- [38] A. Nazir, Um-E-laila, Firdaus-E-bareen, E. Hameed, and M. Shafiq, "Sustainable management of peanut shell through biochar and its application as soil ameliorant," *Sustain.*, vol. 13, no. 24, 2021, doi: 10.3390/su132413796.
- [39] M. S. Kahar, F. Rosalina, I. Febriadi, and A. Fahrizal, "Effect of various extractors and time of shaking on soil reaction and micro element soil," *AIP Conf. Proc.*, vol. 2081, no. March 2019, doi: 10.1063/1.5094025.
- [40] P. D. Johan, O. H. Ahmed, A. Maru, L. Omar, and N. A. Hasbullah, "Optimisation of charcoal and sago (*Metroxylon sago*) bark ash to improve phosphorus availability in acidic soils," *Agronomy*, vol. 11, no. 9, pp. 1–28, 2021, doi: 10.3390/agronomy11091803.
- [41] Z. Liu *et al.*, "Modified biochar: synthesis and mechanism for removal of environmental heavy metals," *Carbon Res.*, vol. 1, no. 1, pp. 1–21, 2022, doi: 10.1007/s44246-022-00007-3.
- [42] D. Firnia, S. Anwar, D. Santosa, B. Nugroho, and D. P. Baskoro, "Transformation of aluminium fractions and phosphorus availability in acid soils as the result of microbes and ameliorant addition," *J. Degrad. Min. L. Manag.*, vol. 7, no. 4, pp. 2355–2362, 2020, doi: 10.15243/jdmlm.
- [43] D. Lauricella *et al.*, "Impact of novel materials on alkalinity movement down acid soil profiles when combined with lime," *J. Soils Sediments*, vol. 21, no. 1, pp. 52–62, 2021, doi: 10.1007/s11368-020-02747-4.
- [44] S. W. Budi, C. Wibowo, A. Sukendro, and H. S. Bektu, "Growth improvement of falcataria moluccana inoculated with mycosilvi grown in post-mining silica sand soil medium amended with soil ameliorants," *Biodiversitas*, vol. 21, no. 1, pp. 422–427, 2020, doi: 10.13057/biodiv/d210149.
- [45] P. Paramisparam, O. H. Ahmed, L. Omar, H. Y. Ch'ng, P. D. Johan, and N. H. Hamidi, "Co-application of charcoal and wood ash to improve potassium availability in tropical mineral acid soils," *Agronomy*, vol. 11, no. 10, pp. 1–30, 2021, doi: 10.3390/agronomy11102081.
- [46] Y. Sun *et al.*, "Roles of biochar-derived dissolved organic matter in soil amendment and environmental remediation: A critical review," *Chem. Eng. J.*, vol. 424, no. April, 2021, doi: 10.1016/j.cej.2021.130387.
- [47] L. Han *et al.*, "Biochar's stability and effect on the content, composition and turnover of soil organic carbon," *Geoderma*, vol. 364, no. January, pp. 1–11, 2020, doi: 10.1016/j.geoderma.2020.114184.
- [48] M. Satria, N. Harun, F. Hamzah, and A. Pramana, "Characteristics of charcoal briquettes corn cobs charcoal with the addition of areca peel charcoal," *J. Phys. Conf. Ser.*, vol. 2049, no. 1, 2021, doi: 10.1088/1742-6596/2049/1/012082.
- [49] Hasanuddin, H. Nurdin, Waskito, and D. Y. Sari, "Characteristic of Areca Fiber Briquettes as Alternative Energy," *J. Phys. Conf. Ser.*, vol. 1594, no. 1, 2020, doi: 10.1088/1742-6596/1594/1/012049.
- [50] N. H. Hamidi, O. H. Ahmed, L. Omar, and H. Y. Ch'ng, "Combined use of charcoal, sago bark ash, and urea mitigate soil acidity and aluminium toxicity," *Agronomy*, vol. 11, no. 9, pp. 1–15, 2021, doi: 10.3390/agronomy11091799.
- [51] P. Paramisparam, O. H. Ahmed, L. Omar, H. Y. Ch'ng, A. Maru, and P. D. Johan, "Amending potassic fertilizer with charcoal and sago (*Metroxylon sago*) bark ash to improve potassium availability in a tropical acid soil," *Agronomy*, vol. 11, no. 11, pp. 1–31, 2021, doi: 10.3390/agronomy11112222.
- [52] P. D. Johan, O. H. Ahmed, L. Omar, and N. A. Hasbullah, "Charcoal and sago bark ash on ph buffering capacity and phosphorus leaching," *Agronomy*, vol. 11, no. 11, pp. 1–20, 2021, doi: 10.3390/agronomy11112223.
- [53] J. Rambli, W. A. W. A. K. Ghani, M. A. M. Salleh, and R. Khezri, "Evaluation of biochar from Sago (*Metroxylon Spp.*) as a potential solid fuel," *BioResources*, vol. 14, no. 1, pp. 1928–1940, 2019, doi: 10.15376/biores.14.1.1928-1940.
- [54] A. B. Syuhada, J. Shamshuddin, C. I. Fauziah, A. B. Rosenani, and A. Arifin, "Biochar as soil amendment: Impact on chemical properties and corn nutrient uptake in a Podzol," *Can. J. Soil Sci.*, vol. 96, no. 4, pp. 400–412, 2016, doi: 10.1139/cjss-2015-0044.
- [55] F. Zulfiqar *et al.*, "Challenges in organic component selection and biochar as an opportunity in potting substrates: a review," *J. Plant Nutr.*, vol. 42, no. 11–12, pp. 1386–1401, 2019, doi: 10.1080/01904167.2019.1617310.
- [56] J. Rawat, J. Saxena, and P. Sanwal, "Biochar: A Sustainable Approach for Improving Plant Growth and Soil Properties," in *Biochar*, V. Abrol, Ed. IntechOpen, 2019, pp. 1–17.
- [57] L. Zhang, Y. Xiang, Y. Jing, and R. Zhang, "Biochar amendment effects on the activities of soil carbon, nitrogen, and phosphorus hydrolytic enzymes: a meta-analysis," *Environ. Sci. Pollut. Res.*, vol. 26, no. 22, 2019, doi: 10.1007/s11356-019-05604-1.
- [58] P. Madhavi, V. Sailaja, T. R. Prakash, and S. Hussain, "Characterization of Biochar and Humic Acid and their Effect on Soil Properties in Maize," *Int. J. Curr. Microbiol. Appl. Sci.*, vol. 6, no. 9, pp. 449–457, 2017, doi: 10.20546/ijcmas.2017.609.054.
- [59] N. Bohari *et al.*, "Nutritional characteristics of biochar from pineapple leaf residue and sago waste," *Pertanika J. Sci. Technol.*, vol. 28, no. Special issue 2, pp. 273–286, 2020, doi: 10.47836/pjst.28.S2.21.
- [60] Y. P. Situmeang, I. D. N. Sudita, and M. Suarta, "Application of Compost and Biochar from Cow, Goat, and Chicken Manure to Restore Soil Fertility and Yield of Red Chili," *Int. J. Adv. Sci. Eng. Inf. Technol.*, vol. 11, no. 5, pp. 2008–2015, 2021, doi: 10.18517/IJASEIT.11.5.13845.
- [61] F. Rosalina, M. A. A. Gafur, I. Irnawati, M. H. Soekamto, Z. Sangadji, and S. M. Kahar, "Utilization of Compost and Zeolite as Ameliorant on Quartz Sand Planting Media for Caisim (*Brassica Juncea*) Plant Growth," *J. Phys. Conf. Ser.*, vol. 1155, no. 1, 2019, doi: 10.1088/1742-6596/1155/1/012055.
- [62] K. Jindo *et al.*, "Role of biochar in promoting circular economy in the agriculture sector. Part 2: A review of the biochar roles in growing media, composting and as soil amendment," *Chem. Biol. Technol. Agric.*, vol. 7, no. 1, pp. 1–10, 2020, doi: 10.1186/s40538-020-00179-3.
- [63] M. R. Nemat, F. Simard, J.-P. Fortin, and J. Beaudoin, "Potential Use of Biochar in Growing Media," *Vadose Zo. J.*, vol. 14, no. 6, p. vzj2014.06.0074, 2015, doi: 10.2136/vzj2014.06.0074.
- [64] Sugiyarto, A. Salim, and R. Firgiyanto, "The effect of the use of various kinds of biochar and soil nutrients on pakcoy (*brassica rapa* l.)," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 672, no. 1, 2021, doi: 10.1088/1755-1315/672/1/012014.
- [65] S. E. Hale *et al.*, "The effect of biochar, lime and ash on maize yield in a long-term field trial in a Ultisol in the humid tropics," *Sci. Total Environ.*, vol. 719, pp. 137455, 2020, doi: 10.1016/j.scitotenv.2020.137455.