













- [32] L. H. Xie *et al.*, "Physicochemical properties of rice starch for production of vermicelli with premium quality," *J. Food Sci. Technol.*, vol. 54, no. 12, pp. 3928–3935, 2017, doi: 10.1007/s13197-017-2852-9.
- [33] I. Sapna and A. Jayadeep, "Application of pulverization and thermal treatment to pigmented broken rice: insight into flour physical, functional and product forming properties," *J. Food Sci. Technol.*, vol. 58, no. 6, pp. 2089–2097, 2021, doi: 10.1007/s13197-020-04718-6.
- [34] P. O. Suklaew, C. Chusak, and S. Adisakwattana, "Physicochemical and functional characteristics of rd43 rice flour and its food application," *Foods*, vol. 9, no. 12, 2020, doi: 10.3390/foods9121912.
- [35] L. M. Fonseca, S. L. M. El Halal, A. R. G. Dias, and E. da R. Zavareze, "Physical modification of starch by heat-moisture treatment and annealing and their applications: A review," *Carbohydr. Polym.*, vol. 274, no. June, p. 118665, 2021, doi: 10.1016/j.carbpol.2021.118665.
- [36] S. Sittipod and Y. C. Shi, "Changes in physicochemical properties of rice starch during steeping in the parboiling process," *J. Cereal Sci.*, vol. 69, pp. 398–405, 2016, doi: 10.1016/j.jcs.2016.05.010.
- [37] K. Kunyane, T. Van Ngo, S. Kusumawardani, and N. Lungsakul, "Ultrasound-chilling assisted annealing treatment to produce a lower glycemic index of white rice grains with different amylose content," *Ultrason. Sonochem.*, vol. 87, no. June, p. 106055, 2022, doi:10.1016/j.ultsonch.2022.106055.
- [38] H. Huang *et al.*, "Comparative quality evaluation of physicochemical and amylose content profiling in rice noodles from diverse rice hybrids in China," *Agric.*, vol. 13, no. 1, 2023, doi:10.3390/agriculture13010140.
- [39] M. Saeri *et al.*, "Formulation of Vermicelli Mixed Corn and Rice Flour with Additional Carrageenan and Its Economic Value," *Int. J. Food Sci.*, vol. 2022, 2022, doi: 10.1155/2022/7387223.
- [40] G. Jacobasch, G. Dongowski, D. Schmiedl, and K. Müller-Schmehl, "Hydrothermal treatment of Novelose 330 results in high yield of resistant starch type 3 with beneficial prebiotic properties and decreased secondary bile acid formation in rats," *Br. J. Nutr.*, vol. 95, no. 6, pp. 1063–1074, Jun. 2006, doi: 10.1079/bjn20061713.
- [41] W. Rungratanawanich, M. Memo, and D. Uberti, "Redox homeostasis and natural dietary compounds: Focusing on antioxidants of rice (*Oryza sativa* L.)," *Nutrients*, vol. 10, no. 11, pp. 1–19, 2018, doi:10.3390/nu10111605.
- [42] S. Yamuangmorn and C. Prom-U-thai, "The potential of high-anthocyanin purple rice as a functional ingredient in human health," *Antioxidants*, vol. 10, no. 6, pp. 1–21, 2021, doi:10.3390/antiox10060833.
- [43] S. B. Sasongko, B. P. Rini, H. Maehiroh, F. D. Utari, and M. Djaeni, "The effect of temperature on vermicelli drying under dehumidified air," *IOP Conf. Ser. Mater. Sci. Eng.*, vol. 1053, no. 1, p. 012102, 2021, doi:10.1088/1757-899x/1053/1/012102.
- [44] S. Yamuangmorn, B. Dell, and C. Prom-U-Thai, "Effects of cooking on anthocyanin concentration and bioactive antioxidant capacity in glutinous and non-glutinous purple rice," *Rice Sci.*, vol. 25, no. 5, pp. 270–278, 2018, doi: 10.1016/j.rsci.2018.04.004.
- [45] T. Laokuldilok and N. Kanha, "Effects of processing conditions on powder properties of black glutinous rice (*Oryza sativa* L.) bran anthocyanins produced by spray drying and freeze drying," *LWT - Food Sci. Technol.*, vol. 64, no. 1, pp. 405–411, 2015, doi:10.1016/j.lwt.2015.05.015.
- [46] M. A. Farooq *et al.*, "Investigating the structural properties and in vitro digestion of rice flours," *Food Sci. Nutr.*, vol. 9, no. 5, pp. 2668–2675, 2021, doi: 10.1002/fsn3.2225.
- [47] L. Wang *et al.*, "Effect of annealing on the physico-chemical properties of rice starch and the quality of rice noodles," *J. Cereal Sci.*, vol. 84, pp. 125–131, 2018, doi: 10.1016/j.jcs.2018.10.004.
- [48] C. F. Li and B. S. Luh, "Rice snack food," in *Rice: Production and utilization*, B. S. Luh, Ed., Westport: Avi Publishing, 1980, pp. 690–711.
- [49] Y. Hu *et al.*, "Influence of dynamic high temperature during grain filling on starch fine structure and functional properties of semi-waxy japonica rice," *J. Cereal Sci.*, vol. 101, no. March, 2021, doi:10.1016/j.jcs.2021.103319.
- [50] J. Singh, L. Kaur, and O. J. McCarthy, "Factors influencing the physico-chemical, morphological, thermal and rheological properties of some chemically modified starches for food applications-A review," *Food Hydrocoll.*, vol. 21, no. 1, pp. 1–22, 2007, doi:10.1016/j.foodhyd.2006.02.006.