











- [9] S. Z. Ramle, S. N. H. Oslan, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Biochemical Characteristics of Acid-Soluble Collagen from Food Processing By-products of Needlefish Skin (*Tylosurus acus melanotus*)," *Applied Sciences*, vol. 12, no. 24, Dec. 2022, doi: 10.3390/app122412695.
- [10] H. Hasanuddin, A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Effect of different acids during collagen extraction the bone and fins from purple-spotted bigeye (*Priacanthus tayenus* Richardson, 1846) and their physicochemical properties," *Food Research*, vol. 8, no. 1, pp. 326–335, Feb. 2024, doi:10.26656/fr.2017.8(1).224.
- [11] A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Extraction and Characterization of Type I Collagen from Parrotfish (*Scarus sordidus* Forsskål, 1775) Scale solubilized with the Aid of Acetic Acid and Pepsin," *International Journal of Biomaterials*, vol. 2023, pp. 1–10, Apr. 2023, doi:10.1155/2023/7312447.
- [12] N. S. Fatiroi, A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Biochemical and Microstructural Characteristics of Collagen Biopolymer from Unicornfish (*Naso reticulatus* Randall, 2001) Bone Prepared with Various Acid Types," *Polymers*, vol. 15, no. 4, Feb. 2023, doi: 10.3390/polym15041054.
- [13] D. F. Harianto, A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, S. M. Rambli, and N. Huda, "Assessing the Organic Acids-solubilized Collagen Derived from the Skin of Unicornfish (*Naso reticulatus* Randall, 2011) on Physicochemical and Structural Characteristics," *ASEAN Journal on Science and Technology for Development*, vol. 40, no. 2, p. 1-12, Feb. 2024, doi: 10.61931/2224-9028.1535.
- [14] G. K. S. Arumugam, D. Sharma, R. M. Balakrishnan, and J. B. P. Ettiyappan, "Extraction, optimization and characterization of collagen from sole fish skin," *Sustainable Chemistry and Pharmacy*, vol. 9, pp. 19–26, Sep. 2018, doi: 10.1016/j.scp.2018.04.003.
- [15] R. Ahmed, M. Haq, and B. S. Chun, "Characterization of marine derived collagen extracted from the by-products of bigeye tuna (*Thunnus obesus*)," *International Journal of Biological Macromolecules*, vol. 135, pp. 668–676, 2019, doi:10.1016/j.ijbiomac.2019.05.213.
- [16] F. Yu, C. Zong, S. Jin, J. Zheng, N. Chen, J. Huang, Y. Chen, F. Huang, Z. Yang, Y. Tang, and G. Ding, "Optimization of extraction conditions and characterization of pepsin-solubilised collagen from skin of giant croaker (*Nibea japonica*)," *Marine Drugs*, vol. 16, no. 1, Jan. 2018, doi: 10.3390/md16010029.
- [17] J. Wang, X. Pei, H. Liu, and D. Zhou, "Extraction and characterization of acid-soluble and pepsin-soluble collagen from skin of loach (*Misgurnus anguillicaudatus*)," *International Journal of Biological Macromolecules*, vol. 106, pp. 544–550, Jan. 2018, doi:10.1016/j.ijbiomac.2017.08.046.
- [18] W. Liu, Y. Zhang, N. Cui, and T. Wang, "Extraction and characterization of pepsin-solubilized collagen from snakehead (*Channa argus*) skin: Effects of hydrogen peroxide pretreatments and pepsin hydrolysis strategies," *Process Biochemistry*, vol. 76, pp. 194–202, Jan. 2019, doi: 10.1016/j.procbio.2018.10.017.
- [19] Department of Fisheries Malaysia, "Fisheries Statistics I 2023," accessed on 22 May 2024, <https://www.dof.gov.my/en/resources/fisheries-statistics-i/>
- [20] A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Wan, and N. Huda, "Tropical Marine Fish Surimi By-products: Utilisation and Potential as Functional Food Application," *Food Review International*, vol. 39, no. 6, p. 3455-3480, Dec. 2021, doi:10.1080/87559129.2021.2012794.
- [21] A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, S. M. Rambli, and N. Huda, "Byproducts of Tropical Marine Processing as a Candidate for Industrial Food Ingredients an Appraisal on Macro and Micronutrients," *ASEAN Journal on Science and Technology for Development*, vol. 40, no. 2, Feb. 2024, doi: 10.61931/2224-9028.1535.
- [22] A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Chemical composition of lizardfish surimi by-product: Focus on macro and micro-minerals contents," *Current Research in Nutrition and Food Science*, vol. 9, no. 1, pp. 52–61, Apr. 2021, doi:10.12944/CRNFSJ.9.1.06.
- [23] A. A. Jaziri, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Physicochemical and Microstructural Analyses of Pepsin-Soluble Collagens Derived from Lizardfish (*Saurida tumbil* Bloch, 1795) Skin, Bone and Scales," *Gels*, vol. 8, no. 8, pp. 1–18, 2022, doi:10.3390/gels8080471.
- [24] N. Huda, E. K. Seow, N. M. Nor, N. A. N. Muhammad, A. Fazilah, and A. M. Easa, "Effect of duck feet collagen addition on physicochemical properties of surimi," *International Food Research Journal*, vol. 20, no. 2, p. 537-544, Oct. 2013.
- [25] A. Jongjareonrak, S. Benjakul, W. Visessanguan, and M. Tanaka, "Skin gelatin from bigeye snapper and brownstripe red snapper: Chemical compositions and effect of microbial transglutaminase on gel properties," *Food Hydrocolloids*, vol. 20, no. 8, pp. 1216–1222, Dec. 2006, doi: 10.1016/j.foodhyd.2006.01.006.
- [26] J. Chen, J. Li, Z. Li, R. Yi, S. Shi, K. Wu, Y. Li, and S. Wu, "Physicochemical and functional properties of type I collagens in red stingray (*Dasyatis akajei*) skin," *Marine Drugs*, vol. 17, no. 10, Sep. 2019, doi: 10.3390/md17100558.
- [27] K. Matmaroh, S. Benjakul, T. Prodpran, A. B. Encarnacion, and H. Kishimura, "Characteristics of acid soluble collagen and pepsin soluble collagen from scale of spotted golden goatfish (*Parupeneus heptacanthus*)," *Food Chemistry*, vol. 129, no. 3, pp. 1179–1186, Dec. 2011, doi: 10.1016/j.foodchem.2011.05.099.
- [28] F. Y. Cheng, F. W. Hsu, H. S. Chang, L. C. Lin, and R. Sakata, "Effect of different acids on the extraction of pepsin-solubilised collagen containing melanin from silky fowl feet," *Food Chemistry*, vol. 113, no. 2, pp. 563–567, Mar. 2009, doi: 10.1016/j.foodchem.2008.08.043.
- [29] M. M. Schmidt, R. C. P. Dornelles, R. Mello, E. H. Kubota, M. Mazutti, A. P. Kempka, and I. M. Demiate, "Collagen extraction process," *International Food Research Journal*, vol. 23, no. 3, p. 913-922, Oct. 2013.
- [30] J. Cao, Q. Duan, X. Liu, X. Shen, and C. Li, "Extraction and Physicochemical Characterization of Pepsin Soluble Collagens from Golden Pompano (*Trachinotus blochii*) Skin and Bone," *Journal of Aquatic Food Product Technology*, vol. 28, no. 8, pp. 837–847, Sep. 2019, doi: 10.1080/10498850.2019.1652216.
- [31] W. K. Song, D. Liu, L. L. Sun, B. F. Li, and H. Hou, "Physicochemical and Biocompatibility Properties of Type I Collagen from the Skin of Nile Tilapia (*Oreochromis niloticus*) for Biomedical Applications," *Mar Drugs*, vol. 17, no. 3, Feb. 2019, doi: 10.3390/md17030137.
- [32] S. N. H. Oslan, R. Shapawi, R. A. M. Mokhtar, W. N. M. Noordin, and N. Huda, "Characterization of Acid- and Pepsin-Soluble Collagen Extracted from the Skin of Purple-Spotted Bigeye Snapper," *Gels*, vol. 8, no. 10, Oct. 2022, doi: 10.3390/gels8100665.
- [33] J. Bakar, U. Hartina, M. Razali, D. Hashim, and A. Q. Sazili, "Properties of collagen from barramundi (*Lates calcarifer*) skin," *International Food Research Journal*, vol. 20, no. 2, p. 835-842, Nov. 2012.
- [34] M. Sadowska, I. Kołodziejska, and C. Niecikowska, "Isolation of collagen from the skins of Baltic cod (*Gadus morhua*)," *Food Chemistry*, vol. 81, no. 2, pp. 257-262, May. 2003, doi:10.1016/S0308-8146(02)00420-X.
- [35] Q. Q. Wu, T. Li, B. Wang, and G. F. Ding, "Preparation and characterization of acid and pepsin-soluble collagens from scales of croceine and redlip croakers," *Food Science and Biotechnology*, vol. 24, no. 6, pp. 2003–2010, Dec. 2015, doi: 10.1007/s10068-015-0264-z.
- [36] Q. Bin Luo, C. F. Chi, F. Yang, Y. Q. Zhao, and B. Wang, "Physicochemical properties of acid- and pepsin-soluble collagens from the cartilage of Siberian sturgeon," *Environmental Science and Pollution Research*, vol. 25, no. 31, pp. 31427–31438, Nov. 2018, doi: 10.1007/s11356-018-3147-z.
- [37] S. Chen, H. Chen, Q. Xie, B. Hong, J. Chen, F. Hua, K. Bai, J. He, R. Yi, and H. Wu, "Rapid isolation of high purity pepsin-soluble type I collagen from scales of red drum fish (*Sciaenops ocellatus*)," *Food Hydrocolloids*, vol. 52, pp. 468–477, Jan. 2016, doi:10.1016/j.foodhyd.2015.07.027.
- [38] S. Iswariya, P. Velswamy, and T. S. Uma, "Isolation and Characterization of Biocompatible Collagen from the Skin of Puffer Fish (*Lagocephalus inermis*)," *J Polym Environ*, vol. 26, no. 5, pp. 2086–2095, May 2018, doi: 10.1007/s10924-017-1107-1.
- [39] M. Atef, S. M. Ojagh, A. M. Latifi, M. Esmacili, and C. C. Udenigwe, "Biochemical and structural characterization of sturgeon fish skin collagen (*Huso huso*)," *Journal of Food Biochemistry*, vol. 44, no. 8, Aug. 2020, doi: 10.1111/jfbc.13256.
- [40] S. Benjakul, Y. Thiansilakul, W. Visessanguan, S. Roytrakul, H. Kishimura, T. Prodpran, and J. Meesane, "Extraction and characterisation of pepsin-solubilised collagens from the skin of bigeye snapper (*Priacanthus tayenus* and *Priacanthus macracanthus*)," *Journal of the Science of Food and Agriculture*, vol. 90, no. 1, pp. 132–138, Jan. 2010, doi: 10.1002/jsfa.3795.

- [41] B. B. Doyle, E. G. Bendit, E. R. Blout, "Infrared Spectroscopy of Collagen and Collagen-Like Polypeptides," *Biopolymers*, vol. 14, no. 5, pp. 937-57, May, 1975, doi: 10.1002/bip.1975.360140505.
- [42] I. Bae, K. Osatomi, A. Yoshida, K. Osako, A. Yamaguchi, and K. Hara, "Biochemical properties of acid-soluble collagens extracted from the skins of underutilised fishes," *Food Chemistry*, vol. 108, no. 1, pp. 49-54, May 2008, doi: 10.1016/j.foodchem.2007.10.039.
- [43] S. Chuaychan, S. Benjakul, and H. Kishimura, "Characteristics of acid- and pepsin-soluble collagens from scale of seabass (*Lates calcarifer*)," *LWT – Food Science and Technology*, vol. 63, no. 1, pp. 71-76, Sep. 2015, doi: 10.1016/j.lwt.2015.03.002.