[29] N. S. Al-Amri, H. A. Ewea, and A. M. Elfeki, “Revisit the rational method for flood estimation in the Saudi arid environment,” Arabian Journal of Geosciences, vol. 15, no. 6, Mar. 2022, doi:10.1007/s12517-021-09219-0.

[30] S. S. Fanta and C. H. Sime, “Performance assessment of SWAT and HEC-HMS model for runoff simulation of Toba watershed, Ethiopia,” Sustainable Water Resources Management, vol. 8, no. 1, Nov. 2021, doi: 10.1007/s40899-021-00596-8.

[31] Al-Amri, Nassir S., Hatem A. Ewea, and Amro M. Elfeki. "Revisit the rational method for flood estimation in the Saudi arid environment." Arabian Journal of Geosciences 15, no. 6 (2022): pp.532.

[32] Pool, Sandra, Marc Vis, and Jan Seibert. "Regionalization for Ungauged Catchments — Lessons Learned From a Comparative Large-Sample Study." Water Resources Research 57, no. 10 (2021): e2021WR030437. Accessed December 8, 2023. doi:10.1029/2021WR030437.

[33] J. Nossent and W. Bauwens, Application of a normalized Nash-Sutcliffe efficiency to improve the accuracy of the Sobol’ sensitivity analysis of a hydrological model. 2012.

[34] Soomro, Abdul Ghani, Muhammad Munir Babar, Anila Hameem Memon, Arjumand Zehra Zaidi, Arshad Ashraf, and Jewell Lund. “Sensitivity of Direct Runoff to Curve Number Using the SCS-CN Method.” Civil Engineering Journal 5, no. 12 (December 1, 2019): 2738–46. doi:10.28991/cej-2019-03091445

[35] Fialová, J., A. M. Negm, and M. Zelenakova. “Introduction to the ‘Assessment and Protection of Water Resources in the Czech Republic.’” Assessment and Protection of Water Resources in the Czech Republic, June 26, 2019, 3–10. doi:10.1007/978-3-030-18363-9_1.

[36] K. R. Douglas-Mankin *et al.*, “A comprehensive review of ephemeral gully erosion models,” *CATENA*, vol. 195, p. 104901, 2020, doi:10.1016/j.catena.2020.104901.

[37] G. Zhang, X. Su, O. O. Ayantobo, K. Feng, and J. Guo, “Spatial interpolation of daily precipitation based on modified ADW method for gauge-scarce mountainous regions: A case study in the Shiyang River Basin,” *Atmospheric Research*, vol. 247, p. 105167, 2021, doi:10.1016/j.atmosres.2020.105167.

[38] A. Arlimasita and U. Lasminto, “Sensitivity analysis of rainfall-runoff model in malino sub-watershed,” International Journal on Advanced Science, Engineering and Information Technology, vol. 10, no. 4, pp. 1578–1583, 2020, doi: 10.18517/ijaseit.10.4.12803.

[39] S. Chebii, M. Mukolwe, and B. Ong’or, “River flow modelling for flood prediction using artificial neural network in ungauged Perkerra catchment, Baringo County, Kenya,” Water Practice and Technology, vol. 17, Apr. 2022, doi: 10.2166/wpt.2022.034.

[40] W. Ben khelifa and M. Mosbahi, “Modeling of rainfall-runoff process using HEC-HMS model for an urban ungauged watershed in Tunisia,” Modeling Earth Systems and Environment, vol. 8, pp. 1–10, Jun. 2022, doi: 10.1007/s40808-021-01177-6.

[41] I. El Ghoul, H. Sellami, S. Khlifi, and M. Vanclooster, “Impact of land use land cover changes on flow uncertainty in Siliana watershed of northwestern Tunisia,” CATENA, vol. 220, p. 106733, 2023, doi:10.1016/j.catena.2022.106733.