



















- 104641, 2022, doi: 10.1016/j.jfca.2022.104641.
- [27] K. B. Beć, J. Grabska, and C. W. Huck, "Miniaturized NIR Spectroscopy in Food Analysis and Quality Control: Promises, Challenges, and Perspectives," *Foods*, vol. 11, no. 10, 2022.
- [28] B. M. Nicolai *et al.*, "Nondestructive measurement of fruit and vegetable quality by means of NIR spectroscopy: A review," *Postharvest Biol. Technol.*, vol. 46, pp. 99–118, 2007, doi:10.1016/j.postharvbio.2007.06.024.
- [29] N. U. Haase, "Prediction of potato processing quality by near infrared reflectance spectroscopy of ground raw tubers," *J. Near Infrared Spectrosc.*, vol. 19, no. 1, pp. 37–45, 2011, doi: 10.1255/jnirs.919.
- [30] L. S. Magwaza *et al.*, "Assessment of rind quality of 'Nules Clementine' mandarin fruit during postharvest storage: 2. Robust Vis/NIRS PLS models for prediction of physico-chemical attributes," *Sci. Hortic. (Amsterdam)*, vol. 165, pp. 421–432, 2014, doi:10.1016/j.scienta.2013.09.050.
- [31] A. Phuphaphud, K. Saengprachatanarug, J. Posom, E. Taira, and L. Panduangnate, "Prediction and Classification of Energy Content in Growing Cane Stalks for Breeding Programmes Using Visible and Shortwave Near Infrared," *Sugar Tech*, vol. 24, no. 5, pp. 1497–1509, 2022, doi: 10.1007/s12355-021-01075-2.
- [32] D. Saputro, D. Priambodo, M. Pahlawan, and R. Masithoh, "Classification of Cocoa Beans Based on Fermentation Level Using PLS-DA Combined with Visible Near-Infrared (VIS-NIR) Spectroscopy," *Proc. 2nd Int. Conf. Smart Innov. Agric. (ICoSIA 2021)*, vol. 19, no. ICoSIA 2021, pp. 100–106, 2022, doi:10.2991/absr.k.220305.015.
- [33] D. N. Rahmia, E. Z. D. Pratiwi, M. F. R. Pahlawan, H. Z. Amanah, and R. E. Masithoh, "Non-destructive measurement of malic acid content of sapodilla fruit using visible near infrared (VisNIR) spectroscopy with variations in storage temperature," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 1116, no. 1, p. 012027, Dec. 2022, doi: 10.1088/1755-1315/1116/1/012027.
- [34] B. Rongtong, T. Suwonsichon, P. Ritthiruangdej, and S. Kasemsumran, "Determination of water activity, total soluble solids and moisture, sucrose, glucose and fructose contents in osmotically dehydrated papaya using near-infrared spectroscopy," *Agric. Nat. Resour.*, vol. 52, no. 6, pp. 557–564, 2018, doi:10.1016/j.anres.2018.11.023.
- [35] R. E. Masithoh, F. Roosmayanti, K. Rismiwandira, and M. F. R. Pahlawan, "Detection of Palm Sugar Adulteration by Fourier Transform Near-Infrared (FT-NIR) and Fourier Transform Infrared (FT-IR) Spectroscopy," *Sugar Tech*, no. Livesey 2003, 2021, doi:10.1007/s12355-021-01058-3.
- [36] B. Hou, Y. Hu, P. Zhang, and L. Hou, "Potato Late Blight Severity and Epidemic Period Prediction Based on Vis/NIR Spectroscopy," *Agric.*, vol. 12, no. 7, 2022, doi: 10.3390/agriculture12070897.
- [37] D. S. Kasampalis, P. Tsouvaltzis, K. Ntouros, A. Gertsis, D. Moshou, and A. S. Siomos, "Rapid nondestructive postharvest potato freshness and cultivar discrimination assessment," *Appl. Sci.*, vol. 11, no. 6, Mar. 2021, doi: 10.3390/app11062630.
- [38] M. Šulc, Z. Kotíková, L. Paznocht, and J. Lachman, "Changes in Carotenoid Profile during Potato (*Solanum tuberosum* L.) Tuber Maturation," *Am. J. Potato Res.*, vol. 98, no. 2, pp. 85–92, 2021, doi:10.1007/s12230-020-09805-0.
- [39] L. M. Londoño-Giraldo, A. M. Baena-Pedroza, F. Martínez-Seidel, E. Corpas-Iguarán, and G. Taborda-Ocampo, "Gone wild: Integration of antioxidative, physicochemical, volatilomic and sensorial profiles ratify rustic relatives of cherry tomato as ideal mating partners," *Sci. Hortic. (Amsterdam)*, vol. 277, no. October 2020, 2021, doi:10.1016/j.scienta.2020.109814.